



Structural and Functional Changes in Periodontal Tissues During Prosthetics with Metal-Ceramic and Zirconium Dentures

1. Axmedov Xurshid Kamalovich

Received 2nd Oct 2023,
Accepted 19th Nov 2023,
Online 27th Dec 2023

¹ Bukhara State Medical Institute

Abstract: To increase the efficiency of prosthetics, the use of metal, metal-ceramic and zircon dentures was determined. The issues of modern methods of dental prosthetics are highlighted. The analysis of the most common errors and complications in prosthetics with the use of fixed structures has been carried out. The assessment of the physical and technical properties of cermet structures remains poorly studied. To exclude ceramic chips, antagonizing dentures are not brought to contact, which can lead to deformation. Thus, the question of the influence of fixed prostheses on the clinical and morphological state in the absence of periodontal tissues remains unstudied.

Introduction

In case of prosthetics with a non-removable design of dentures, special importance is attached to the abutment teeth. They carry out a large functional load, in connection with which the periodontal tissue undergoes morpho-functional changes. Considering the side effects of artificial crowns, these changes become obvious. Initial structural changes are observed in the pulp tissue during the preparation of dental hard tissues. As stated by Professor Kh.I. Irsaliev during the preparation of the teeth, the temperature of the pulp of the tooth rises due to the high speed of the turbine drill. Electron microscopic observations indicate blood filling, and in some cases hemorrhage of the dental pulp, and this contributes to aseptic inflammation of the dental pulp tissues. At the same time, the choice of design features primarily depends on the state and degree of changes in the periodontal tissues. According to certain data, pronounced morpho-functional changes are observed in the periodontal microvessels with an increase in functional loads. It should be borne in mind that the problems associated with obtaining reliable information on the condition of the tissues surrounding the tooth (periodontium, periodontium, marginal gum) arise already at the initial stage. The diagnostic methods used are rather approximate and in many cases do not reflect the actual picture of the state of the periodontal tissues.

Thus, when preparing hard tissues of teeth and design features of fixed structures of dentures with various materials of artificial crowns, they contribute to profound changes in the pathological nature of the periodontal tissues.

In orthopedic dentistry, dentures from precious and base metal alloys are widely used to restore defects in crowns and dentition. The latter are composed of cobalt-chrome and nickel-chrome alloys. The problem of the influence of base metals on the organs of the oral cavity and on the body as a whole is urgent. Among them, oral galvanosis is more common. Being in the oral cavity for a long time, a denture, as a foreign body made of a material not characteristic of the human body, causes structural changes in the marginal periodontium. Therefore, modern orthopedic dentistry faces the question of the compatibility of metal dental alloys and the human body [1.3.5.7.9.11.13.15.17].

The phenomenon of "intolerance", galvanosyndrome, galvanosis described in the literature, can be explained by the body's reaction to the presence of metallic inclusions in the oral cavity, which is expressed not only by intolerance to dentures, but also by the manifestation of a burning sensation of the tongue, hypo- and hypersalivation, changes in taste, etc. etc. In this regard, the study of the state of galvanosis of the oral cavity with fixed prosthetics made of various metals becomes urgent and requires further research in this direction, which will improve the quality of dental care and prevent premature removal of orthopedic structures. The available research in this direction is not numerous and is of a fragmentary nature. Therefore, the identification of the cause-and-effect circumstances that contribute to the occurrence and cause of diseases of oral mucosa when using various types of prostheses (metal crowns, ceramics, zirconium) with the use of histological, electron microscopic and histochemical research methods, will allow developing more effective methods of their treatment and prevention.

Materials and methods

The studies were carried out in 100 people, who were divided into 2 groups: the main observation group - 90 ($80.7 \pm 3.7\%$) patients and the comparison group (control group) - 30 ($19.3 \pm 3.7\%$) patients. Among the patients of the main group, there are 30 patients with chromium-cobalt, 30 patients with metal-ceramics and 30 patients with zircon crowns. There were examined 144 teeth, which are and will be the abutment teeth of bridges. In the comparison group, studies of 24 teeth with intact periodontium on the upper and lower jaws were carried out in patients aged 30 to 60 years. When examining all patients in both observation groups, the Green-Vermillion hygienic index, Russell periodontal index were determined, and the Kulazhenko test was performed. The Schiller-Pisarev test refers to the method of intravital staining of gingival glycogen, the content of which increases with chronic inflammation (composition - crystalline iodine 1 g, potassium iodide 2 g, distilled water 40 ml). Intense discoloration after lubrication of the gums indicates inflammation. Oral cavity sanitation, dental plaque removal, and oral hygiene training were carried out. Diagnostic observations of the periodontal condition were carried out in the area of abutment teeth before and after fixation of bridges and crowns, as well as within 1, 3, 6, 12 months after prosthetics. Clinical methods included questioning, casual examination, and armed eye examination (stomatoscopy) of the oral cavity. Stomatoscopy was performed using an ophthalmic operating microscope at a magnification of 30 times.

For morphological examination, 1×1.5 mm tissue pieces were taken for transmission electron microscopy. Pieces of bioplast were fixed in glutaraldehyde. Electron microscopic studies were carried out under the guidance of Professor I.M. Baybekov. The bone tissue around the abutment tooth was examined using X-ray machines. The pictures were taken according to Yusupov. Computed tomography was performed to assess the bone structure of the upper and lower jaw. To determine the endurance of the periodontal tissue to vertical load, gnathodynamometry was performed using electronic gnathodynamometers. Electroodontometry was performed selectively, and only those teeth that were included in the abutment teeth for bridges and according to indications. The used bridges (cermets, zirconium, metal crowns) create conditions for the development of increased functional load on the periodontal tissues of the supporting teeth. When prosthetics of included dentition defects with

bridges, it is necessary to determine possible functional changes in the periodontal tissues, which can largely determine the terms of their use. The change in blood circulation in the periodontal tissues directly depends on the degree of deformation of the dentition, the length of the edentulous and the involvement of bridges in the chewing load. Particular attention during prosthetics (cermets, zirconium) should be paid to changes in the state of the marginal periodontium at the level of microcirculation, which was one of the objectives of this study. To determine the marginal periodontium of the abutment teeth of the first group, we selected 64 patients, the second group - 10 people.

The study groups consisted of 5 groups: 10 patients ($16.0 \pm 4.2\%$) - with intact periodontal disease, 16 patients ($24.0 \pm 4.9\%$) - gingivitis, 22 patients ($36.0 \pm 5.5\%$) - mild periodontitis, 10 patients ($13.3 \pm 3.9\%$) - periodontitis of moderate severity, 6 patients ($10.7 \pm 3.5\%$) - periodontitis at the stage of abscess formation. The state of the periodontal tissues, in particular the marginal gums, was assessed according to the data of clinical and paraclinical research methods in the field of abutment teeth in the dynamics of observations up to 1 year in patients of the main observation group and the comparison group. Statistical processing of the obtained data was carried out using the variation statistics of the computer programs "MS Excell" and "MS Access" in a standard volume for biomedical research. Before using the technique of electroodontometry, dental deposits were removed. During the second prosthetics, after removing the bridges, the remains of the fixed cement were carefully removed. EOM, intraoral image. The study of the EOM of teeth with a filling in the area of the tooth neck was not carried out in order to avoid the current leaving the soft tissues. The existing amalgam fillings were removed and measurements were taken from the bottom of the carious cavity. Patients with contact fillings on the teeth underwent preliminary isolation of the teeth with a celluloid plate smeared with petroleum jelly, after which the indicators were measured. Electroodontometry was carried out using an EOM-3 apparatus.

Immunological studies were carried out in the laboratory of immunoregulation of the institute of the Immunology and Human Genomics of Academy of Sciences of the Republic of Uzbekistan. The study involved 180 patients (115 women and 65 men), aged 20 to 70 years old. The average age of women was 38 years old, men - 49 years old. The patients were conditionally divided into two groups:

1st group - 180 patients (115 women and 65 men) who underwent metal-ceramic prosthetics;

Group 2 - 30 healthy patients (17 women and 13 men) who underwent zirconium prosthetics.

The control was the data of a survey of 30 apparently healthy volunteers, comparable in age and sex.

Measurement of concentration of IL-1 β , IL-6, IL-10, TNF α in oral liquid (GC), was conducted by enzyme immunoassay (ELISA) using commercial kits AO "Vector-Best". Gastrointestinal tract sampling was performed before complex treatment in patients with defects in dentition and hard dental tissues and after prosthetics, not earlier than after 30 days, as well as in healthy individuals. To do this, at the dental appointment, the patient was asked to rinse the oral cavity with 3 small volumes of plain water and think about lemon to stimulate salivation. Sterile penicillin vials with a volume of 10 ml were used as test tubes, which patients filled with their gastric cancer up to 1 / 3-1 / 2 of the total volume. Then glass vials with the test material were frozen in a freezer and stored at minus 20 ° C for no more than one month. Thereafter, undiluted samples were thawed very quickly by heat treatment in a water bath at 37 ° C to prevent fibrinogen precipitation.

The data were statistically processed using the Statistics 6.0 computer program. The reliability of differences in the mean values of the compared indicators was assessed by the Student's test (t).

Clinical researches

The collection of clinical material was carried out with the obligatory opening of an outpatient card, entering into it the dental status and the general condition of the body and the diseases transferred. All design features and type of material from which bridges were made, age indicators are given in the section "Material and research methods". In the comparison group (control group), studies of 24 teeth with intact periodontium in the upper and lower jaws were carried out on the basis of basic and paraclinical research methods (percussion, the state of the gingival margin using a bulbous probe), X-ray - overview and sighting images, electroodontometry - to determine condition of the pulp and periodontal. In the main observation group of patients, in addition to determining the type of design features of bridges, all abutment teeth were examined in the same sequence as in the comparison group.

All patients were familiarized with the treatment and manipulation plan (removal of bridges if necessary, the forthcoming stages of treatment, replacement of constructions, excluding possible side and pathological conditions), written consent and financial costs signed by patients [2.4.6.8.10.12.14.16.18.20].

Criteria for the quality of crowns were assessed by the inconsistency of anatomical shapes, taking into account the group of teeth, excessive or insufficient insertion of crowns into the gingival groove, causing retraction of the tooth neck or leading to hypertrophic gingivitis, supercontacts - to changes in the periodontal complex. Taking into account the radiological, functional parameters and the state of the periodontal complex, electroodontometry (EOM) was used: the abutment teeth were subjected to depulpation or replanting of the root canals with poor-quality obturation. During prosthetics, the abutment teeth were subjected to odontopreparation under cermet or zirconium according to the generally accepted technique: the design of the ledge, retraction of the periodontal edge using the appropriate threads. The impression was obtained using a silicone sandwich technique. Laboratory and clinical stages: fitting of frames, prosthesis, fixation was carried out according to the rules of orthopedic dentistry.

The clinical condition of the marginal periodontium was assessed visually and using an operating microscope, the Schiller-Pisarev test and the papillary-marginal alveolar index (PMA) indexes, where hyperemia, edema, bleeding and vascular pattern were determined.

The clinical picture with partial adentia largely depends on the number of lost teeth, dystopia, type of bite, functional usefulness, the state of hard tissues not only of abutment teeth, but also of the complex of tissues surrounding the tooth. The appearance of a defect after tooth extraction disrupts its continuity, morphological and functional disorders occur. Shortened periods of wearing fixed bridges resulted in supercontact, while in others the teeth were turned off from the act of chewing. The marginal gum, where the connection with the tooth occurs, is often interpreted as a gingival pocket, a gingival gap, a physiological pocket, a gingival groove. When measuring the gingival gap, the groove depth reached 2-3 mm on average. With bridge prosthetics, when odontopreparation was performed and the gums were injured (periodontal pathology), the depth of the gingival groove in 50% of cases reached 3 mm. These data related to the indicators of the marginal gingiva of the abutment teeth around the defect during repeated prosthetics of patients. The condition of the periodontal gap served as a basis for assessing the marginal periodontium in persons wearing fixed prostheses, both during re-prosthetics and in future orthopedic interventions. We studied the magnitude of the electrochemical potentials in connection with the finding in the oral cavity of prostheses of various design features, as well as the materials from which they were made: solid, cermets, zircon [17.19.21.23.24].

It is known that dentures made of various materials lead to disruption of oral homeostasis and have adverse effects on the body as a whole. In particular, there are violations in the triggering mechanisms

in the development of allergies, hyperergic conditions, ulcerative and trophic processes. In this regard, the determination and understanding of causal factors in the conditions of the galvanic environment of the oral cavity is based on the use of para-clinical methods, in particular, on the study of electrochemical potentials. Our data on electrochemical potentials (ECP) are consonant with the indicators of numerous studies and are normally up to 50 mV. Analyzing the identified ECP frequencies, it is obvious that wearing metal structures is directly related to an increase in the level of galvanic currents. Thus, when studying ECP in patients using structures made of dissimilar materials, the average measurement values tended to higher values of the potential difference and reached the level of 100-150 mV. Poor, inadequate endodontic treatment leads to changes in the periodontal complex, ultimately leading to side effects and premature removal of bridges.

When removing bridges from metal brazed-stamped structures, it turned out that in 95% of cases, dark-colored corrosion, covered with nutritive masses, were observed at the places of the rations of the prosthesis body with supporting crowns, and under the prosthesis body there were bedsores of the mucous membrane of bright red color, hypertrophy, bleeding. The errors observed by us in the work of a dentist and a technician at the stages of manufacturing fixed dentures were assessed by the following parameters: modeling of the manufacture of crowns and the intermediate part of the prosthesis body; supercontacts, which, if incorrect, leads to abrasion of the teeth of the antagonists and abrasion of the facing material. The above data of clinical and paraclinical known techniques for fixed bridge prosthetics indicate the presence of pathological conditions that entail the premature removal of bridges.

We evaluated the periodontal complex in 90 patients who received orthopedic treatment. Of these, 38 patients underwent primary and 20 patients underwent repeated orthopedic treatment with fixed bridges, with the complete exclusion of acrylic, plastic veneers. We examined 114 teeth under the supports of bridges and 24 teeth on the upper and lower jaw with healthy intact periodontium in patients of the control group. The dynamics of observations began with planning and coordinating with the patients the design features of bridges and the upcoming therapeutic measures in the dynamics of treatment up to 1 year. Selectively, in terms of up to 3 years and more, after prosthetics, some patients were called for a follow-up examination, where, if necessary, a clinical and X-ray examination was carried out. In patients using for many years (more than 7 years) a fixed construction of dentures made of cobalt alloy coated with titanium sodium, an unpleasant odor from the oral cavity was noted. The mucous membrane of the gums is hyperemic, sometimes bluish in color, repeats the shape of the neck of artificial crowns. Bleed easily and painful on probing. In some patients (more in men), the titanium coating has worn off on the chewing surfaces of the teeth. Patients with such (metal) dental constructions complained of the appearance of a metallic taste in the mouth and the sensation of unpleasant micro-currents when using aluminum spoons. In the dynamics of observations in terms of up to 1 year and more with repeated prosthetics, we paid special attention to the state of the periodontal edge, the mucous membrane of the entire oral cavity, modeling in accordance with the anatomy of the group of teeth, supercontacts, the state of the intermediate part of the bridge, as well as functional and aesthetic indicators.

Organo-structural changes both in the pulp of the tooth and periodontium, where electrodontometry showed increased indicators: the current strength reached up to 100-150 μ A (at a rate of 2 to 5 μ A), which unambiguously predetermined the study of the state of the surrounding tooth of the marginal gum, taking into account the common genesis ... In connection with the significant prevalence of diseases of the oral mucosa caused by a reaction to the materials of dentures, histological and histochemical studies of the oral mucosa using prostheses made of various materials are of great interest. The structural and functional characteristics of the mucous membrane of the marginal gum is of great applied and practical importance, as it allows one to judge the correct choice of materials for

dentures, which in the process of being in the oral cavity may be indifferent, excluding possible pathological conditions in the future. In this regard, one should take into account the histological picture of the mucosa, which can become an essential guideline for determining the pathological nature of the oral mucosa with various prostheses.

The aim of this study is the histological and histochemical assessment of the oral mucosa with various types of prostheses: cermets, zirconium, metal crowns. The control group consisted of marginal gum tissue after tooth extraction. The keratinization process is especially pronounced in the epithelium of the anterior third of the hard palate. The structure of the epithelium of the hard palate is very similar to the leaves that have fallen to the ground in the autumn period of the year. Some of them slough off in the form of completely keratinized scales, while others do not show any signs of keratinization and have a structure typical of squamous epithelium. Regeneration of epithelial cells occurs due to cell division of the basal layers. The pattern of mitosis, as a rule, is observed in the cells of the basal layer, as well as in the adjacent lower rows of cells of the spiny layer. (Irsaliev Kh.I. 1993). The electron diffraction pattern of a transmission microscope shows detachment of cells of the granular layer with penetration of microorganisms into the intercellular space. Destructive changes are also observed in the deep zones of the spiny layers of the epithelium. It should be noted a slowdown in the differentiation of epithelial cells and a sharp decrease in the mitotic activity of the combined zone. In the prickly layers, there is a rupture of desmosomic connections between epithelial cells.

With the introduction of metals (mainly stainless steel) into the practice of orthopedic treatment, the reactions to these materials were explained by the appearance of galvanic phenomena in the oral cavity. As a rule, the diagnosis was formulated as "galvanosis", "electrogalvanosis". Since there were no sufficiently sensitive research methods to prove the presence of metal corrosion in an aggressive environment. Significant disorders with partial edentulousness occur in the tissues of the periodontal complex, since it is the integrity of the dentition of the jaws that ensures the usefulness of chewing as one of the most important functions of the body. The wide spread of secondary adentia among the adult population makes this problem relevant all over the world Gozhiy A.G. 1998, Volyntsev V.I. 1996). In the manufacture of a fixed structure of dentures, most of the bridges have to be prepared with abutment teeth for artificial crowns. Dissection of hard dental tissues without observing the established rules leads to morpho-functional changes in dental pulp tissues. In the monograph of Professor Irsaliev Kh.I. (2001) widely described and illustrated electronograms of the surface of prepared teeth.

The author describes the structure of the paired tooth surface using burs of different dispersion under a scanning electron microscope. Coarsely dispersed burs form deep uneven reliefs of the tooth surface, which lead to burnout and dentin necrosis of the teeth (12,15). The author considers burs with fine-grained diamond coatings to be the most acceptable for preparation. Experimental and clinical studies have shown that when prosthetics of included dentition defects with bridges, it is necessary to carefully determine the indications for the use of various structures, taking into account the reactive changes in the periodontal tissues. The functional state of periodontal tissues is largely determined by their trophism, which is provided by the circulatory system.

In the works of Arslanov O.U. "Compensatory and adaptive processes in the dentition with partial secondary adentia and the ways of their orthopedic correction" (2007), a sharp violation of microcirculation and a decrease in the endurance of periodontal tissues of abutment teeth was proved, which, according to the author, is a contraindication for immediate prosthetics after the removal of decayed teeth or their roots (20).

Laser Doppler flowmetry is a method for diagnosing and monitoring trophic changes in periodontal tissues and during orthopedic interventions to give an idea of the hemodynamic state of the corresponding area. When studying the functional state of periodontal vessels in secondary adentia, many authors have established changes in the marginal gingiva and abnormalities in the pulp of

abutment teeth after their preparation (21,22,23). Hemodynamic changes in the microcirculation of the supporting teeth of the marginal gum against the background of single studies cannot give an objective assessment that reflects the true picture when wearing or long-term use of bridges. In our opinion, special attention should be paid to the measurement of the state of the marginal periodontium in modern types of prosthetics (cermets, metal, zirconium) at the level of microcirculation, adaptation processes, which was the topic of this study.

We studied the state of the marginal periodontium of abutment teeth in 70 people aged 20 to 60 years, who underwent laser Doppler flowmetry as part of the research. Dissection of teeth (depulped and with live pulp) was carried out in a strict sequence: creating a ledge, reducing the length of the tooth crown taking into account the ceramic and metal layer, removing hard tooth tissues from the vestibular and oral surfaces, and forming a circular ledge at the level of the gingival margin. The work used diamond burs in a certain sequence, taking into account the formation of the ledge. After taking impressions with antagonists with a silicone mass, the teeth were covered with temporary crowns, which were pre-fabricated by taking an impression and making them with plastic. Subsequently, prosthetics were carried out according to the generally accepted technique: fitting a solid frame, choosing a color, eliminating supercontacts, fixing a bridge with glass ionomer cement.

Results and discussion

Thus, the restoration of dentition defects with various fixed structures of dentures is an orthopedic dental treatment. The loss of teeth and the appearance of various defects in the dentition contribute to the further development of the pathological process in the dento-jaw system and in the body as a whole. The choice of the design of dentures and the materials from which they will be made is of paramount importance in the medical and social sphere of the patient. In our Republic of Uzbekistan and a number of other foreign countries, metal crowns and bridges made of base metals are currently used, sometimes they are coated with titanium nitrate, which further aggravates the protective mechanisms of the oral cavity.

Such materials used for dentures do not meet either the aesthetic or functional requirements of modern dentistry. This state of the problem dictated the search for the most adequate structures built into the oral cavity, devoid of the above disadvantages. Currently, the most satisfying clinical and functional requirements and aesthetic parameters are dentures made of cermet, zirconium. Undoubtedly, the positive results of orthopedic structures made of cermets and zirconium are largely indifferent for the organs and tissues of the oral cavity. However, there are also negative influences on the marginal periodontium associated with the advancement of the edges of the crowns into the periodontal pocket, causing morphological changes. The present study is devoted to the study of the marginal periodontium when using fixed bridges, cermets, zirconium and chromium-cobalt alloy).

As recommendations for practical health care of the most appropriate optimal designs, it is necessary to use cermets, zirconium in order to improve the quality of dental care. In patients using bridges and crowns made of a cobalt-chromium alloy, edematous with cyanosis of the mucous membrane of the alveolar ridge was observed. As we mentioned above, such a mucous membrane quickly bleeds when touched with a probe and when brushing teeth with a toothbrush.

We also described structural changes in the epithelial cover at the level of light and electron microscopy. We associate all these changes with the appearance of galvanic currents in the mouth in patients. Determination of electrochemical potentials in the mouth in these patients showed the maximum values and were equal to 120-150 mV. A high level of galvanic currents in the oral cavity, exceeding the physiological norm, dictates the implementation of preventive measures to prevent the occurrence of galvanism in the oral cavity, which should include the selection of design features that are tolerant to the physiological homeostasis of the oral cavity, compliance with the complete homogeneity of metal structures and their replacement in the presence of heterogeneity of inclusions.

In this regard, the study of the functional state of microcirculation in the tissues of the marginal gums and the entire periodontal complex is relevant for obtaining further positive results of prosthetics of dentition defects. In this regard, when choosing a plan for orthopedic treatment, the assessment of the functional state of the supporting-retaining apparatus of the teeth comes to the fore. However, problems arise in the first approximation and are associated with an objective assessment of the true picture of the state of the tissues surrounding the tooth. The currently used diagnostic methods are quite subjective: there are no standard measurement techniques, developed protocols, and the available data are contradictory. Thus, the existing methods for assessing the state of the marginal gingiva of the periodontal abutment teeth do not give a true picture of its functional changes in microcirculation at the stages of prosthetics with secondary adentia.

In our opinion, special attention should be paid to determining the state of the marginal periodontium of abutment teeth in modern types of prosthetics (cermets, zirconium), adaptation processes at the level of microcirculation, which was the topic of the study. We have examined by clinical, functional and morphological methods the tissues of the marginal periodontium of abutment teeth in patients using metal-ceramic and zircon bridges.

It should be noted that at present there is no multifactorial approach to the diagnosis, treatment and prevention of diseases caused by the materials of dentures, which significantly complicates the practical work of a dentist. Therefore, the histological diagnosis of the mucous membrane of the marginal periodontium will certainly contribute to the differential diagnosis of diseases caused by the materials of dentures. The development of inflammatory diseases is determined by the state of cytokine regulation. Most of both pro- and anti-inflammatory cytokines are present not only in the peripheral blood, but also in other biological fluids of the body [9]. The sources of their production are both lymphocytes and macrophages built into the epithelium of the mucous membranes, and the epithelial cells of the mucous membranes and the salivary glands themselves. Another source of cytokines in gastric cancer (saliva) may be their extravasation from blood serum. However, many researchers have noted that the content of cytokines in gastric cancer (saliva) does not correlate with their level in the blood, which indirectly indicates their local synthesis.

As a rule, the content of FN O- α is not determined, or is at a low level in the blood serum of healthy people, whereas with the development of a pathological process, its amount increases several times. Thus, synthesis FN O- α in patients with the group to prosthetic averaged 12.3 ± 0.32 pg / ml, and these values were in the control group 9, 4 ± 0.39 pg / ml, which is 1, 3 times more ($P < 0.001$). It was also found that a significant increase in the level of IL-10 in the main observation group before orthodontic treatment was 1.2 times (10.6 ± 0.32 pg / ml versus 8.5 ± 0.38 pg / ml) ($P < 0.001$).

The obtained data with cytokine status of the oral cavity before the prosthetics point to the weakening of the local immune defense GPRS, connectivity with the need for the orthopedic intervention.

After the imposition of dentures, the 1st dental examination was carried out the next day, at the stage of denture correction. Treatment of inflammatory changes in the mucous membrane was carried out both at the reception and at home using preparations containing antiseptic, anesthetic, regenerating components in accordance with the "Protocol for the management of patients with partial absence of teeth (partial secondary adentia)".

Thus, in the I-group Ceramic prostheses after 7 days, 83% (72) patient s complained of pain in the oral mucosa during interdigitation and upon application of the denture. Examination revealed inflammatory changes in the mucous membrane of the prosthetic bed, which were represented by hyperemia and erosion, localized in the region of the border of the denture, as well as in areas of increased base

pressure on the mucous membrane. At the repeated stages of the denture correction, the patients showed no complaints, no inflammatory changes were found on the oral mucosa.

The study of the content of cytokines showed that in patients with reddening of the mucous membrane, the area of which coincided with the shape of the edge of the metal-ceramic denture, already on the 7th day, a cytokine imbalance was noted in the oral fluid, characterized by a significant increase in the content of IL-1 β , IL-6, IL-10, and FN O- α (Fig. 1).

Comparative analysis of the content revealed that in group I with metal-ceramic prosthetics, the level of IL-1 β increased by 3 times after a week with an average value of 22.7 ± 0.35 pg / ml (before treatment, 7.5 ± 0.20 pg / ml) ($P < 0.001$), the concentration of IL-6 increased to 42.9 ± 1.07 pg / ml (before treatment 14.7 ± 0.46 pg / ml) ($P < 0.001$), TNF- α - 37.2 ± 0.76 pg / ml (before treatment 12.3 ± 0.32 pg / ml) ($P < 0.001$), IL-10 - 21.5 ± 0.55 pg / ml (before treatment 10.6 ± 0.32 pg / ml) ($P < 0.001$) (Fig. 1).

Comparative analysis of the obtained data of group II with zirconium prosthetics revealed a positive dynamics of indices of local immunity links in gastric cancer in patients of this group. On examination, no foci of inflammation caused by the prostheses were observed, which indicates a satisfactory regeneration of the tissues of the prosthetic bed, and according to the patients, they were completely adapted to the denture.

The obtained data of the comparative analysis make it possible to develop an algorithm for the treatment of the consequences of metal-ceramic and zirconium prostheses, in the use of orthopedic structures made of various materials.

Conclusions

Dental prosthetics in patients using chromium-cobalt pure metal bridges with a coating of titanium dioxide) have a negative effect on the marginal periodontium and the oral mucosa, manifested by high values of electrochemical potentials, unsatisfactory hygienic state, profound changes in the morphological structures of the marginal periodontal complex in the dentoparodontium. Laser Doppler flowmetry at the stages of orthopedic treatment showed that the functional state of the microcirculation system in the tissues of the marginal gums of the periodontal tissues is relevant for predicting the results of prosthetics of partial dentition defects.

The structural and functional characteristics of the mucous membrane of the marginal gingiva indicated the negative effect of metal dentures in comparison with the weakly expressed inflammation of the epithelium in the surface layers during prosthetics made of metal-ceramics and zirconium. Diagnostics and treatment of an increased level of galvanic currents should involve the removal of metal structures of various inclusions and the conduct of pathogenetic therapy of periodontal tissues before re-prosthetics, with the replacement of basic dental materials with ceramics or zirconium.

The activation of the inflammatory process in the periodontal tissues was established, manifested by an increase in the concentration of pro- and anti-inflammatory cytokines already on the 7th day after fixation of the prosthetic structure. It was revealed that with zirconium prosthetics, the positive dynamics of indices of local immunity links in gastric cancer in patients of this group. On examination, no foci of inflammation caused by the prostheses were observed, which indicates a satisfactory regeneration of the tissues of the prosthetic bed, and in the patients' opinion, they were completely adapted to the denture.

REFERENCES

1. Х.И.Ирсадиев Сканирующая электронная микроскопия твердых тканей зубов при их патологической стираемости. Ж. Stomatologiya 2002, №3-4 стр.16-21

2. Ирсалиев Х.И., Рахмонов Х.Ш. Структурная организация тканей зубов при их патологиях. Ж. Stomatologiya 2003, № 1-2, стр.110-116
3. Ражабов О.А., Иноятлов А.Ш. Металлокерамика ва циркон протезлардан фойдаланилганда оғиз бўшлиғидаги клиник функционал ўзгаришларнинг диагностика тахлиллари //Услубий тавсиянома. – Бухоро - 2021
4. Ирсалиев Х.И., Зуфаров С.А., Мавляноходжаев И., Байбеков И.М. Морфологические особенности взаимодействия при сменочной микрофлоры с эпителиоцитами слизистой оболочки альвеолярного гребня у больных с частичным отсутствием зубов. Журнал Стоматология. 1991, №6 стр. 48-50
5. Ирсалиев Х.И. Ультроструктурные особенности слизистой оболочки альвеолярного гребня при частичном отсутствии зубов. Медицинский журнал Узбекистана, 1985 №9 стр.82
6. Капитонова Ю.М. Лечение вторичной адентии мостовидными протезами // Автореф. дис.канд.мед.наук, Москва, 2002.
7. Байбеков И.М., Ирсалиев Х.И., Мавляноходжаев И. Взаимодействие индигенных пристеночных микроорганизмов с клетками слизистой оболочки пищеварительного тракта. Архив патологии, 1992 №5 стр.18-24
8. Байбеков И.М., Ирсалиев Х.И., Мавляноходжаев И. Ультраструктурные аспекты взаимодействия микроорганизмов с клетками слизистой оболочки пищеварительного тракта. 14 Всесоюзная конференция по электронной микроскопии. Москва 1992, стр.78
9. Хабилов Н.Л., Байбеков И.М., Ирсалиев Х.И. Морфологическая оценка биосовместимости различных компонентов биоситама. Медицинский журнал Узбекистана, 2003, № 3 стр.74-77
10. Брагин Е.А. Тактика зубодесневого сохранения при протезировании несъемными протезами // Стоматология 2003;4:44-47.
11. Волынец В.М. Клинико-лабораторные показатели непереносимости сплавов металлов // Автореф. дисс. канд. мед.наук. Киев, 1996.- с. 17.
12. Ирсалиев Х.И., Байбеков И.М., Ханазаров Д.А., Рахманов Х.Ш. Функциональная морфология барьерно-защитных комплексов полости рта. Монография 2001, Ташкент
13. Арутюнов С.Д. Профилактика осложнений при применении металлокерамических зубных протезов // Автореф. дис. канд. мед.наук. М 1990.- 19с.
14. Банченко Г.В. Флейшер Г.М. и др. Электроманнитная аллергия-гальваноз Медицинский алфавит // Стоматология, 2/2012.- с. 42-51.
15. Ермолов В.В. Лазерная профилактика пародонтальных осложнений при ортопедическом лечении несъемными протезами // Автореф. дисс. канд. мед.наук, Москва, 2004.
16. Ирсалиев Х.И. Особенности барьерно-защитных комплексов полости рта до и в процессе пользования зубными протезами. Автор.дисс.доктор.мед.наук 1993 стр.17
17. Гожий А.Г. Клинические проявления электрохимических процессов, обусловленных отделочной обработкой зубных протезов из нержавеющей стали/ А.Г. Гожий, Г.Р. Сагатеян, Л.Д. Гожая // Стоматология. – 1998. –№3.- с. 46-50.
18. Ирсалиев Х.И. с соавт. Сканирующая электронная микроскопия зубов после одонтопрепарирования и покрытия Fluorprotector. Журнал Стоматология 2000, №4 стр.25-28

19. Арсланов О.У. Компенсаторно-приспособительные процессы в зубо-челюстной системе при частичной и вторичной адентии и пути их ортопедической коррекции. // Автореф. дисс. канд. мед.наук, Ташкент 2007
20. Ирсалиев Х.И., Гафуров Г.А. Восстановление дефектов зубных рядов мостовидными протезами и их влияние на гемодинамику тканей пародонта. Ж. Stomatologiya № 3-4, 2006 (33-34) с.24-26
21. Ирсалиев Х.И., Сафаров М.Т., Хасанова Н. Гемодинамические изменения слизистой оболочки альвеолярного гребня нижней челюсти при частичных дефектах зубных рядов. Ж. Stomatologiya № 1-2, 2010 (41-42) стр. 79-85.
22. Бякова Ж.С. Оценка состояния регионарного кровотока при использовании зубных протезов // Нижегород.Мед.Журн., 2003.- с.164-165.
23. Ражабов О.А. Клинические изменения полости рта при использовании металлокерамических зубных протезов// Ж.Тиббиётдаянги кун 1(29)2020 стр.322-325.
24. Rajabov O.A. Clinical and functional changes the oral cavity using ceramic-metal dentures//South Asian Academic Research Journals Vol.10 issue2, February2020.209-215.

